Index of Authors

ABD EL MONEIM, A. M. *see* CHRISTIANSEN, S.
ACOSTA-GALLEGOS, J. A. *see* CASTELLANOS, J. Z.
AGRICULTURAL RESEARCH MODELLERS' GROUP, Proceedings of the Twenty-Seventh Meeting of the Agricultural Research Modellers' Group, 117
AINSLEY, A. E. *see* HIDE, G. A.
AL-ASMI, H. S. *see* ESECHIE, H. A.
AMIN, MD R. and THOMAS, H. Growth and water relations of diverse populations of *Lolium perenne* exposed to drought in field, glasshouse and controlled environment, 15
AZAM-ALI, S. N. *see* COLLINSON, S. T.

BADAWI, M. A. *see* EL-SAYED, S. F.
BAINBRIDGE, A. *see* JENKYN, J. F.
BARRY, T. N. *see* SEMIADI, G., WANG, Y.
BIRTLES, M. J. *see* HOCKING EDWARDS, J. E.
BOATMAN, N. D. *see* GRUNDY, A. C.
BORTOLUSSI, G., TERNOUTH, J. H. and MCMENIMAN, N. P. Dietary nitrogen and phosphorus depletion in cattle and their effects on liveweight gain, blood metabolite concentrations and phosphorus kinetics, 493
BORTOLUSSI, G. *see also* TERNOUTH, J. H.
BOUJENNA, A. *see* RAMOS, J. M.
BROCK, J. L., HUME, D. E. and FLETCHER, R. H. Seasonal variation in the morphology of perennial ryegrass (*Lolium perenne*) and cocksfoot (*Dactylis glomerata*) plants and populations in pastures under intensive sheep grazing, 37
BROOKMAN, S. K. E. *see* TALLOWIN, J. R. B.
BROWN, J., DALE, M. F. B. and MACKAY, G. R. General adaptability of potato genotypes selected in the UK for the Mediterranean region, 441

CASTELLANOS, J. Z., PEÑA-CABRIALES, J. J. and ACOSTA-GALLEGOS, J. A. ^{15}N-determined dinitrogen fixation capacity of common bean (*Phaseolus vulgaris*) cultivars under water stress, 327
CHALK, P. M. *see* SEN, S.
CHAVULA, K. M. *see* COLLINSON, S. T.
CHRISTENSEN, B. T. *see* THOMSEN, I. K.
CHRISTIANSEN, S., ABD EL MONEIM, A. M., COCKS, P. S. and SINGH, M. Seed yield and hardseededness of two amphicarpic pasture legumes (*Vicia sativa* ssp. *amphicarpa* and *Lathyrus ciliolatus*) and two annual medics (*Medicago rigidula* and *M. noeana*), 421
COATES, D. B. *see* TERNOUTH, J. H.
COCKS, P. S. *see* CHRISTIANSEN, S., PORQUEDDU, C.
COLLINSON, S. T., AZAM-ALI, S. N., CHAVULA, K. M. and HODSON, D. A. Growth, development and yield of bambara groundnut (*Vigna subterranea*) in response to soil moisture, 307
CONE, J. W., VAN GELDER, A. H. and MARVIN, H. J. P. Influence of drying method and ageing on chemical and physical properties and *in vitro* degradation characteristics of grass and maize samples, 7

DALE, M. F. B. *see* BROWN, J.
DAVIES, D. B., GARWOOD, T. W. D. and ROCHFORD, A. D. H. Factors affecting nitrate leaching from a calcareous loam in East Anglia, 75
DELAUNE, R. D. *see* SAHRAWAT, K. L.
DIATTA, S. *see* SAHRAWAT, K. L.
DIVERIO, S., GODDARD, P. J. and GORDON, I. J. Physiological responses of farmed red deer to management practices and their modulation by long-acting neuroleptics, 211
DOUGLAS, G. B. *see* WANG, Y.
DYKE, G. V. *see* JENKYN, J. F.

ELIAS, S. *see* ESECHIE, H. A.
EL-SAYED, S. F., BADAWI, M. A., MAHMOUD, I. M. and MOTAWA, M. H. Some morphological characters and yield of broad bean (*Vicia faba*) as affected by pesticide treatments, 463
ERSKINE, W. Seed-size effects on lentil (*Lens culinaris*) yield potential and adaptation to temperature and rainfall in West Asia, 335
ESCOTT-WATSON, P. L. *see* MARAIS, J. P.
ESECHIE, H. A., ELIAS, S., RODRIGUEZ, V. and AL-ASMI, H. S. Response of sunflower (*Helianthus annuus*) to planting pattern and population density in a desert climate, 455
EVANS, D. R., HUMPHREYS, M. O. and WILLIAMS, T. A. Forage yield and quality interactions between white clover and contrasting ryegrass varieties in grazed swards, 295

FAN, L. *see* WANG, X. W.
FIGENSCHOU, D. L. *see* MARAIS, J. P.
FLETCHER, R. H. *see* BROCK, J. L.
FOOTE, A. G. *see* WANG, Y.
FORD, M. A. *see* MOOT, D. J.
FREITAS, B. M. and PAXTON, R. J. The role of wind and insects in cashew (*Anacardium occidentale*) pollination in NE Brazil, 319
FROUD-WILLIAMS, R. J. *see* GRUNDY, A. C.

GARCÍA DEL MORAL, L. F. *see* RAMOS, J. M.
GARWOOD, T. W. D. *see* DAVIES, D. B.
GODDARD, P. J. *see* DIVERIO, S.
GOODEN, J. M. *see* NEUTZE, S. A.
GORDON, I. J. *see* DIVERIO, S.
GRUNDY, A. C., BOATMAN, N. D. and FROUD-WILLIAMS, R. J. Effects of herbicide and nitrogen fertilizer application on grain yield and quality of wheat and barley, 379
GUTIÉRREZ BOEM, F. H. and LAVADO, R. S. The effects of soil sodicity on emergence, growth, development and yield of oilseed rape (*Brassica napus*), 169

HARRIS, P. M. *see* HOCKING EDWARDS, J. E.
HENDERSON, A. L. *see* MOOT, D. J.

HENDRICKSEN, R. E. *see* TERNOUTH, J. H.
HENRIKSEN, K. *see* RUBÆK, G. H.
HIDE, G. A., WELHAM, S. J., READ, P. J. and AINSLEY, A. E. The yield of potato plants as affected by stem canker (*Rhizoctonia solani*), blackleg (*Erwinia carotovora* subsp. *atroseptica*) and by neighbouring plants, 429
HIRANO, C. *see* OWUSU, E. O.
HOCKING EDWARDS, J. E., BIRTLES, M. J., HARRIS, P. M., PARRY, A. L., PATERSON, E., WICKHAM, G. A. and MCCUTCHEON, S. N. Pre- and post-natal wool follicle development and density in sheep of five genotypes, 363
HODSON, D. A. *see* COLLINSON, S. T.
HOLMES, C. W. *see* SEMIADI, G.
HORIIKE, M. *see* OWUSU, E. O.
HUCKLE, C. A. *see* ROOK, A. J.
HUME, D. E. *see* BROCK, J. L.
HUMPHREYS, M. O. *see* EVANS, D. R.

INSA, J. A. *see* RAMOS, J. M.

JAMIESON, P. D. *see* MOOT, D. J.
JENKYN, J. F., TODD, A. D., BAINBRIDGE, A. and DYKE, G. V. Effects of inoculum sources on the accuracy and precision of experiments testing different times of applying fungicides to control powdery mildew (*Erysiphe graminis* f. sp. *hordei*) on spring barley, 259
JONES, M. P. *see* SAHRAWAT, K. L.

KIM, C.-S. *see* OWUSU, E. O.
KNOTT, C. M. The effect of cover crops on the establishment and yield of vining peas and on nitrate leaching, 471

LAI, J. R. *see* WANG, X. W.
LAVADO, R. S. *see* GUTIÉRREZ BOEM, F. H.
LINNEMANN, A. R. Phenological development in bambara groundnut (*Vigna subterranea*) transferred from 14 to 11 h photoperiods, 175
LOI, A. *see* PORQUEDDU, C.

MCCUTCHEON, S. N. *see* HOCKING EDWARDS, J. E.
MCGRATH, S. P. and ZHAO, F. J. Sulphur uptake, yield responses and the interactions between nitrogen and sulphur in winter oilseed rape (*Brassica napus*), 53
MACKAY, G. R. *see* BROWN, J.
MCLEAN, R. W. *see* TERNOUTH, J. H.
MCMENIMAN, N. P. *see* BORTOLUSSI, G.
MAHMOUD, I. M. *see* EL-SAYED, S. F.
MARAIS, J. P., FIGENSCHOU, D. L., ESCOTT-WATSON, P. L. and WEBBER, L. N. Administration in suspension-form of n-alkane external markers for dry matter intake and diet selection studies, 207
MARVIN, H. J. P. *see* CONE, J. W.
MISSELBROOK, T. H., SHEPHERD, M. A. and PAIN, B. F. Sewage sludge applications to grassland: influence of sludge type, time and method of application on nitrate leaching and herbage yield, 343
MIYAMOTO, S. *see* SUTOH, M.
MOOT, D. J., JAMIESON, P. D., HENDERSON, A. L., FORD, M. A. and PORTER, J. R. Rate of change in harvest index during grain-filling of wheat, 387
MOSELEY, G. *see* MTENGETI, E. J., WILMAN, D.
MOTAWA, M. H. *see* EL-SAYED, S. F.
MTENGETI, E. J., WILMAN, D. and MOSELEY, G. Differences between twelve forage species in physical breakdown when eaten, 287
MTENGETI, E. J. *see* WILMAN, D.
MUIR, P. D. *see* SEMIADI, G.
MULBAH, C. K. *see* SAHRAWAT, K. L.

NEUTZE, S. A., GOODEN, J. M. and ODDY, V. H. Uptake of labelled phenylalanine into different blood fractions in the portal vein and cranial mesenteric vein of lambs, 511
NSAHLAI, I. V. and UMUNNA, N. N. Comparison between reconstituted sheep faeces and rumen fluid inocula and between *in vitro* and *in sacco* digestibility methods as predictors of intake and *in vivo* digestibility, 235
NTARE, B. R. *see* SIVAKUMAR, M. V. K.

OBARA, Y. *see* ROSE, M. T., SUTOH, M.
ODDY, V. H. *see* NEUTZE, S. A.
ORPHANOS, P. I. Direct and residual effect of phosphorus on dryland barley, 137
OWEN, E. *see* PIKE, D. J.
OWUSU, E. O., KIM, C.-S., HORIIKE, M. and HIRANO, C. Comparative biological and enzymatic studies on some host-adapted populations of melon and cotton aphid, *Aphis gossypii* (Homoptera: Aphididae), 449

PAIN, B. F. *see* MISSELBROOK, T. H.
PARRY, A. L. *see* HOCKING EDWARDS, J. E.
PATERSON, E. *see* HOCKING EDWARDS, J. E.
PATRICK, W. H. JR *see* SAHRAWAT, K. L.
PAXTON, R. J. *see* FREITAS, B. M.
PEÑA-CABRIALES, J. J. *see* CASTELLANOS, J. Z.
PETERSEN, J. *see* RUBÆK, G. H.
PIKE, D. J., OWEN, E. and SAID, A. N. The dacron bag technique for comparing rumen degradability of untreated and ammonia-treated barley straw – effect of particle size and degree of replication, 201
PILBEAM, C. J. Variation in harvest index of maize (*Zea mays*) and common bean (*Phaseolus vulgaris*) grown in a marginal rainfall area of Kenya, 1
PORQUEDDU, C., LOI, A. and COCKS, P. S. Hardseededness and pattern of hard seed breakdown in Sardinian populations of *Medicago polymorpha* under field conditions, 161
PORTER, J. R. *see* MOOT, D. J.
PURCHAS, R. W. *see* WANG, Y.

RAMOS, J. M., GARCÍA DEL MORAL, L. F., BOUJENNA, A., SERRA, J., INSA, J. A. and ROYO, C. Grain yield, biomass and leaf area of triticale in response to sowing date and cutting stage in three contrasting Mediterranean environments, 253
RASMUSSEN, B. *see* RUBÆK, G. H.
READ, P. J. *see* HIDE, G. A.
ROBERTS, J. M. *see* SIVAKUMAR, M. V. K.
ROCHFORD, A. D. H. *see* DAVIES, D. B.
RODRIGUEZ, V. *see* ESECHIE, H. A.
ROOK, A. J. and HUCKLE, C. A. Sources of variation in the grazing behaviour of dairy cows, 227
ROSE, M. T. and OBARA, Y. Effect of growth hormone on the response to insulin and glucose turnover in sheep, 107
ROYO, C. *see* RAMOS, J. M.

RUBÆK, G. H., HENRIKSEN, K., PETERSEN, J., RASMUSSEN, B. and SOMMER, S. G. Effects of application technique and anaerobic digestion on gaseous nitrogen loss from animal slurry applied to ryegrass (*Lolium perenne*), 481

SAHRAWAT, K. L., MULBAH, C. K., DIATTA, S., DELAUNE, R. D., PATRICK, W. H. JR, SINGH, B. N. and JONES, M. P. The role of tolerant genotypes and plant nutrients in the management of iron toxicity in lowland rice, 143

SAID, A. N. *see* PIKE, D. J.

SEMIADI, G., HOLMES, C. W., BARRY, T. N. and MUIR, P. D. Effects of cold conditions on heat production by young sambar (*Cervus unicolor*) and red deer (*Cervus elaphus*), 221

SEN, S. and CHALK, P. M. Stimulation of root growth and soil nitrogen uptake by foliar application of urea to wheat and sunflower, 127

SERRA, J. *see* RAMOS, J. M.

SHEPHERD, M. A and SYLVESTER-BRADLEY, R. Effect of nitrogen fertilizer applied to winter oilseed rape (*Brassica napus*) on soil mineral nitrogen after harvest and on the response of a succeeding crop of winter wheat to nitrogen fertilizer, 63

SHEPHERD, M. A. *see* MISSELBROOK, T. H.

SINGH, B. N. *see* SAHRAWAT, K. L.

SINGH, M. *see* CHRISTIANSEN, S.

SIVAKUMAR, M. V. K., NTARE, B. R. and ROBERTS, J. M. Growth, yield and plant–water relations of four cowpea (*Vigna unguiculata*) cultivars in the Sahel, 183

SLAFER, G. A. Differences in phasic development rate amongst wheat cultivars independent of responses to photoperiod and vernalization. A viewpoint of the intrinsic earliness hypothesis, 403

SOMMER, S. G. *see* RUBÆK, G. H.

STOUT, D. G. *see* THOMPSON, D. J.

STÜR, W. W. *see* WONG, C. C.

SUTOH, M., OBARA, Y. and MIYAMOTO, S. The effect of sucrose supplementation on kinetics of nitrogen, ruminal propionate and plasma glucose in sheep, 99

SYLVESTER-BRADLEY, R. *see* SHEPHERD, M. A

TALLOWIN, J. R. B. and BROOKMAN, S. K. E. The impact of differences in nitrogen content, nitrogen utilization and loss from laminae on competition between four grass species in an old pasture, 25

TERNOUTH, J. H., BORTOLUSSI, G., COATES, D. B., HENDRICKSEN, R. E. and MCLEAN, R. W. The phosphorus requirements of growing cattle consuming forage diets, 503

TERNOUTH, J. H. *see also* BORTOLUSSI, G.

THOMAS, H. *see* AMIN, MD R.

THOMPSON, D. J. and STOUT, D. G. Influence of sowing rate on dry matter yield, plant density and survival of lucerne (*Medicago sativa*) under dryland and irrigated conditions, 301

THOMSEN, I. K. and CHRISTENSEN, B. T. Availability to subsequent crops and leaching of nitrogen in ^{15}N-labelled sugarbeet tops and oilseed rape residues, 191

TODD, A. D. *see* JENKYN, J. F.

UMUNNA, N. N. *see* NSAHLAI, I. V.

VAN GELDER, A. H. *see* CONE, J. W.

WAGHORN, G. C. *see* WANG, Y.

WANG, X. W., LAI, J. R., FAN, L. and ZHANG, R. B. Effects of recurrent selection on populations of various generations in wheat by using the Tai Gu single dominant male-sterile gene, 397

WANG, Y., DOUGLAS, G. B., WAGHORN, G. C., BARRY, T. N. and FOOTE, A. G. Effect of condensed tannins in *Lotus corniculatus* upon lactation performance in ewes, 353

WANG, Y., DOUGLAS, G. B., WAGHORN, G. C., BARRY, T. N., FOOTE, A. G. and PURCHAS, R. W. Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*), 87

WEBBER, L. N. *see* MARAIS, J. P.

WELHAM, S. J. *see* HIDE, G. A.

WICKHAM, G. A. *see* HOCKING EDWARDS, J. E.

WILLIAMS, T. A. *see* EVANS, D. R.

WILMAN, D., MTENGETI, E. J. and MOSELEY, G. Physical structure of twelve forage species in relation to rate of intake by sheep, 277

WILMAN, D. *see also* MTENGETI, E. J.

WONG, C. C. and STÜR, W. W. Persistence of tropical forage grasses in shaded environments, 151

ZHANG, R. B. *see* WANG, X. W.

ZHAO, F. J. *see* MCGRATH, S. P.

Index of Subjects

Amino acids
Uptake of labelled phenylalanine into different blood fractions in the portal vein and cranial mesenteric vein of lambs (Neutze *et al.*), 511
Anacardium
The role of wind and insects in cashew (*Anacardium occidentale*) pollination in NE Brazil (Freitas and Paxton), 319

Bambara groundnut, *see Vigna*
Barley
Direct and residual effect of phosphorus on dryland barley (Orphanos), 137
The dacron bag technique for comparing rumen degradability of untreated and ammonia-treated barley straw – effect of particle size and degree of replication (Pike *et al.*), 201
Effects of inoculum sources on the accuracy and precision of experiments testing different times of applying fungicides to control powdery mildew (*Erysiphe graminis* f. sp. *hordei*) on spring barley (Jenkyn *et al.*), 259
Effects of herbicide and nitrogen fertilizer application on grain yield and quality of wheat and barley (Grundy *et al.*), 379
***Brassica* crops**
Sulphur uptake, yield responses and the interactions between nitrogen and sulphur in winter oilseed rape (*Brassica napus*) (McGrath and Zhao), 53
Effect of nitrogen fertilizer applied to winter oilseed rape (*Brassica napus*) on soil mineral nitrogen after harvest and on the response of a succeeding crop of winter wheat to nitrogen fertilizer (Shepherd and Sylvester-Bradley), 63
The effects of soil sodicity on emergence, growth, development and yield of oilseed rape (*Brassica napus*) (Gutiérrez Boem and Lavado), 169
Availability to subsequent crops and leaching of nitrogen in ^{15}N-labelled sugarbeet tops and oilseed rape residues (Thomsen and Christensen), 191
Breeding
General adaptability of potato genotypes selected in the UK for the Mediterranean region (Brown *et al.*), 441
Broad bean, *see Vicia faba*

Cashew, *see Anacardium*
Cattle
Sources of variation in the grazing behaviour of dairy cows (Rook and Huckle), 227
Dietary nitrogen and phosphorus depletion in cattle and their effects on liveweight gain, blood metabolite concentrations and phosphorus kinetics (Bortolussi *et al.*), 493
The phosphorus requirements of growing cattle consuming forage diets (Ternouth *et al.*), 503
Climate
Seed-size effects on lentil (*Lens culinaris*) yield potential and adaptation to temperature and rainfall in West Asia (Erskine), 335
Common bean, *see Phaseolus*
Cotton
Comparative biological and enzymatic studies on some host-adapted populations of melon and cotton aphid, *Aphis gossypii* (Homoptera: Aphididae) (Owusu *et al.*), 449
Cowpea, *see Vigna*

Daylength
Phenological development in bambara groundnut (*Vigna subterranea*) transferred from 14 to 11 h photoperiods (Linnemann), 175
Deer
Physiological responses of farmed red deer to management practices and their modulation by long-acting neuroleptics (Diverio *et al.*), 211
Effects of cold conditions on heat production by young sambar (*Cervus unicolor*) and red deer (*Cervus elaphus*) (Semiadi *et al.*), 221
Diets
Administration in suspension-form of n-alkane external markers for dry matter intake and diet selection studies (Marais *et al.*), 207
Digestibility
Influence of drying method and ageing on chemical and physical properties and *in vitro* degradation characteristics of grass and maize samples (Cone *et al.*), 7
Comparison between reconstituted sheep faeces and rumen fluid inocula and between *in vitro* and *in sacco* digestibility methods as predictors of intake and *in vivo* digestibility (Nsahlai and Umunna), 235
Differences between twelve forage species in physical breakdown when eaten (Mtengeti *et al.*), 287
Drought
Variation in harvest index of maize (*Zea mays*) and common bean (*Phaseolus vulgaris*) grown in a marginal rainfall area of Kenya (Pilbeam), 1
Growth and water relations of diverse populations of *Lolium perenne* exposed to drought in field, glasshouse and controlled environment (Amin and Thomas), 15
Growth, yield and plant–water relations of four cowpea (*Vigna unguiculata*) cultivars in the Sahel (Sivakumar *et al.*), 183

Influence of sowing rate on dry matter yield, plant density and survival of lucerne (*Medicago sativa*) under dryland and irrigated conditions (Thompson and Stout), 301
^{15}N-determined dinitrogen fixation capacity of common bean (*Phaseolus vulgaris*) cultivars under water stress (Castellanos *et al.*), 327
Response of sunflower (*Helianthus annuus*) to planting pattern and population density in a desert climate (Esechie *et al.*), 455

Feed intake
Administration in suspension-form of n-alkane external markers for dry matter intake and diet selection studies (Marais *et al.*), 207
Comparison between reconstituted sheep faeces and rumen fluid inocula and between *in vitro* and *in sacco* digestibility methods as predictors of intake and *in vivo* digestibility (Nsahlai and Umunna), 235
Physical structure of twelve forage species in relation to rate of intake by sheep (Wilman *et al.*), 277

Forage
Influence of drying method and ageing on chemical and physical properties and *in vitro* degradation characteristics of grass and maize samples (Cone *et al.*), 7
Persistence of tropical forage grasses in shaded environments (Wong and Stür), 151
Physical structure of twelve forage species in relation to rate of intake by sheep (Wilman *et al.*), 277
Differences between twelve forage species in physical breakdown when eaten (Mtengeti *et al.*), 287
The phosphorus requirements of growing cattle consuming forage diets (Ternouth *et al.*), 503

Genetics
Effects of recurrent selection on populations of various generations in wheat by using the Tai Gu single dominant male-sterile gene (Wang *et al.*), 397

Germination
Hardseededness and pattern of hard seed breakdown in Sardinian populations of *Medicago polymorpha* under field conditions (Porqueddu *et al.*), 161
Seed yield and hardseededness of two amphicarpic pasture legumes (*Vicia sativa* ssp. *amphicarpa* and *Lathyrus ciliolatus*) and two annual medics (*Medicago rigidula* and *M. noeana*) (Christiansen *et al.*), 421

Glucose
The effect of sucrose supplementation on kinetics of nitrogen, ruminal propionate and plasma glucose in sheep (Sutoh *et al.*), 99
Effect of growth hormone on the response to insulin and glucose turnover in sheep (Rose and Obara), 107

Grasses
Influence of drying method and ageing on chemical and physical properties and *in vitro* degradation characteristics of grass and maize samples (Cone *et al.*), 7
The impact of differences in nitrogen content, nitrogen utilization and loss from laminae on competition between four grass species in an old pasture (Tallowin and Brookman), 25
Persistence of tropical forage grasses in shaded environments (Wong and Stür), 151

Grazing
Seasonal variation in the morphology of perennial ryegrass (*Lolium perenne*) and cocksfoot (*Dactylis glomerata*) plants and populations in pastures under intensive sheep grazing (Brock *et al.*), 37
Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*) (Wang *et al.*), 87
Sources of variation in the grazing behaviour of dairy cows (Rook and Huckle), 227
Physical structure of twelve forage species in relation to rate of intake by sheep (Wilman *et al.*), 277
Forage yield and quality interactions between white clover and contrasting ryegrass varieties in grazed swards (Evans *et al.*), 295

Growth rate
Differences in phasic development rate amongst wheat cultivars independent of responses to photoperiod and vernalization. A viewpoint of the intrinsic earliness hypothesis (Slafer), 403

Harvest index
Variation in harvest index of maize (*Zea mays*) and common bean (*Phaseolus vulgaris*) grown in a marginal rainfall area of Kenya (Pilbeam), 1
Rate of change in harvest index during grain-filling of wheat (Moot *et al.*), 387

Helianthus
Stimulation of root growth and soil nitrogen uptake by foliar application of urea to wheat and sunflower (Sen and Chalk), 127
Response of sunflower (*Helianthus annuus*) to planting pattern and population density in a desert climate (Esechie *et al.*), 455

Hormones
Effect of growth hormone on the response to insulin and glucose turnover in sheep (Rose and Obara), 107

Insects
The role of wind and insects in cashew (*Anacardium occidentale*) pollination in NE Brazil (Freitas and Paxton), 319
Comparative biological and enzymatic studies on some host-adapted populations of melon and cotton aphid, *Aphis gossypii* (Homoptera: Aphididae) (Owusu *et al.*), 449

Intercropping
Variation in harvest index of maize (*Zea mays*) and common bean (*Phaseolus vulgaris*) grown in a marginal rainfall area of Kenya (Pilbeam), 1

Iron
The role of tolerant genotypes and plant nutrients in the management of iron toxicity in lowland rice (Sahrawat *et al.*), 143

Irrigation
Influence of sowing rate on dry matter yield, plant density and survival of lucerne (*Medicago sativa*) under dryland and irrigated conditions (Thompson and Stout), 301
Growth, development and yield of bambara groundnut (*Vigna subterranea*) in response to soil moisture (Collinson *et al.*), 307

Lactation
Effect of condensed tannins in *Lotus corniculatus* upon lactation performance in ewes (Wang *et al.*), 353
Leaching
Factors affecting nitrate leaching from a calcareous loam in East Anglia (Davies *et al.*), 75
Availability to subsequent crops and leaching of nitrogen in ^{15}N-labelled sugarbeet tops and oilseed rape residues (Thomsen and Christensen), 191
Sewage sludge applications to grassland: influence of sludge type, time and method of application on nitrate leaching and herbage yield (Misselbrook *et al.*), 343
The effect of cover crops on the establishment and yield of vining peas and on nitrate leaching (Knott), 471
Legumes
Seed yield and hardseededness of two amphicarpic pasture legumes (*Vicia sativa* ssp. *amphicarpa* and *Lathyrus ciliolatus*) and two annual medics (*Medicago rigidula* and *M. noeana*) (Christiansen *et al.*), 421
Lens
Seed-size effects on lentil (*Lens culinaris*) yield potential and adaptation to temperature and rainfall in West Asia (Erskine), 335
Lentil, *see* Lens
Lolium
Growth and water relations of diverse populations of *Lolium perenne* exposed to drought in field, glasshouse and controlled environment (Amin and Thomas), 15
Seasonal variation in the morphology of perennial ryegrass (*Lolium perenne*) and cocksfoot (*Dactylis glomerata*) plants and populations in pastures under intensive sheep grazing (Brock *et al.*), 37
Forage yield and quality interactions between white clover and contrasting ryegrass varieties in grazed swards (Evans *et al.*), 295
Effects of application technique and anaerobic digestion on gaseous nitrogen loss from animal slurry applied to ryegrass (*Lolium perenne*) (Rubæk *et al.*), 481
Lotus
Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*) (Wang *et al.*), 87
Lotus
Effect of condensed tannins in *Lotus corniculatus* upon lactation performance in ewes (Wang *et al.*), 353
Lucerne, *see Medicago*

Maize
Variation in harvest index of maize (*Zea mays*) and common bean (*Phaseolus vulgaris*) grown in a marginal rainfall area of Kenya (Pilbeam), 1
Influence of drying method and ageing on chemical and physical properties and *in vitro* degradation characteristics of grass and maize samples (Cone *et al.*), 7
Medic, *see* Medicago
Medicago
Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*) (Wang *et al.*), 87
Hardseededness and pattern of hard seed breakdown in Sardinian populations of *Medicago polymorpha* under field conditions (Porqueddu *et al.*), 161
Influence of sowing rate on dry matter yield, plant density and survival of lucerne (*Medicago sativa*) under dryland and irrigated conditions (Thompson and Stout), 301
Seed yield and hardseededness of two amphicarpic pasture legumes (*Vicia sativa* ssp. *amphicarpa* and *Lathyrus ciliolatus*) and two annual medics (*Medicago rigidula* and *M. noeana*) (Christiansen *et al.*), 421
Metabolism
The effect of sucrose supplementation on kinetics of nitrogen, ruminal propionate and plasma glucose in sheep (Sutoh *et al.*), 99
Effect of growth hormone on the response to insulin and glucose turnover in sheep (Rose and Obara), 107
Dietary nitrogen and phosphorus depletion in cattle and their effects on liveweight gain, blood metabolite concentrations and phosphorus kinetics (Bortolussi *et al.*), 493
Mixed pastures
Seasonal variation in the morphology of perennial ryegrass (*Lolium perenne*) and cocksfoot (*Dactylis glomerata*) plants and populations in pastures under intensive sheep grazing (Brock *et al.*), 37
Forage yield and quality interactions between white clover and contrasting ryegrass varieties in grazed swards (Evans *et al.*), 295
Models
Proceedings of the Twenty-Seventh Meeting of the Agricultural Research Modellers' Group (Agricultural Research Modellers' Group), 117
Sources of variation in the grazing behaviour of dairy cows (Rook and Huckle), 227
Effects of inoculum sources on the accuracy and precision of experiments testing different times of applying fungicides to control powdery mildew (*Erysiphe graminis* f. sp. *hordei*) on spring barley (Jenkyn *et al.*), 259
Growth, development and yield of bambara groundnut (*Vigna subterranea*) in response to soil moisture (Collinson *et al.*), 307
Rate of change in harvest index during grain-filling of wheat (Moot *et al.*), 387
Differences in phasic development rate amongst wheat cultivars independent of responses to photoperiod and vernalization. A viewpoint of the intrinsic earliness hypothesis (Slafer), 403
The yield of potato plants as affected by stem canker (*Rhizoctonia solani*), blackleg (*Erwinia carotovora* subsp. *atroseptica*) and by neighbouring plants (Hide *et al.*), 429

Nitrogen
The impact of differences in nitrogen content, nitrogen utilization and loss from laminae on competition between four grass species in an old pasture (Tallowin and Brookman), 25
Sulphur uptake, yield responses and the interactions between nitrogen and sulphur in winter oilseed rape (*Brassica napus*) (McGrath and Zhao), 53
Effect of nitrogen fertilizer applied to winter oilseed rape (*Brassica napus*) on soil mineral nitrogen after harvest and on the response of a succeeding crop of winter wheat to nitrogen fertilizer (Shepherd and Sylvester-Bradley), 63

Factors affecting nitrate leaching from a calcareous loam in East Anglia (Davies *et al.*), 75
Stimulation of root growth and soil nitrogen uptake by foliar application of urea to wheat and sunflower (Sen and Chalk), 127
Availability to subsequent crops and leaching of nitrogen in ^{15}N-labelled sugarbeet tops and oilseed rape residues (Thomsen and Christensen), 191
The effect of cover crops on the establishment and yield of vining peas and on nitrate leaching (Knott), 471
Dietary nitrogen and phosphorus depletion in cattle and their effects on liveweight gain, blood metabolite concentrations and phosphorus kinetics (Bortolussi *et al.*), 493

Nitrogen fertilizers
Effect of nitrogen fertilizer applied to winter oilseed rape (*Brassica napus*) on soil mineral nitrogen after harvest and on the response of a succeeding crop of winter wheat to nitrogen fertilizer (Shepherd and Sylvester-Bradley), 63
Effects of herbicide and nitrogen fertilizer application on grain yield and quality of wheat and barley (Grundy *et al.*), 379

Nitrogen fixation
^{15}N-determined dinitrogen fixation capacity of common bean (*Phaseolus vulgaris*) cultivars under water stress (Castellanos *et al.*), 327

Oilseed rape, *see Brassica* crops

Pastures
The impact of differences in nitrogen content, nitrogen utilization and loss from laminae on competition between four grass species in an old pasture (Tallowin and Brookman), 25
Sewage sludge applications to grassland: influence of sludge type, time and method of application on nitrate leaching and herbage yield (Misselbrook *et al.*), 343

Peas, *see Pisum*

Pesticides
Effects of inoculum sources on the accuracy and precision of experiments testing different times of applying fungicides to control powdery mildew (*Erysiphe graminis* f. sp. *hordei*) on spring barley (Jenkyn *et al.*), 259
Effects of herbicide and nitrogen fertilizer application on grain yield and quality of wheat and barley (Grundy *et al.*), 379
Some morphological characters and yield of broad bean (*Vicia faba*) as affected by pesticide treatments (El-Sayed *et al.*), 463

Pests and diseases
Effects of inoculum sources on the accuracy and precision of experiments testing different times of applying fungicides to control powdery mildew (*Erysiphe graminis* f. sp. *hordei*) on spring barley (Jenkyn *et al.*), 259
The yield of potato plants as affected by stem canker (*Rhizoctonia solani*), blackleg (*Erwinia carotovora* subsp. *atroseptica*) and by neighbouring plants (Hide *et al.*), 429
Comparative biological and enzymatic studies on some host-adapted populations of melon and cotton aphid, *Aphis gossypii* (Homoptera: Aphididae) (Owusu *et al.*), 449

Phaseolus
Variation in harvest index of maize (*Zea mays*) and common bean (*Phaseolus vulgaris*) grown in a marginal rainfall area of Kenya (Pilbeam), 1
^{15}N-determined dinitrogen fixation capacity of common bean (*Phaseolus vulgaris*) cultivars under water stress (Castellanos *et al.*), 327

Phosphorus
Direct and residual effect of phosphorus on dryland barley (Orphanos), 137
Dietary nitrogen and phosphorus depletion in cattle and their effects on liveweight gain, blood metabolite concentrations and phosphorus kinetics (Bortolussi *et al.*), 493
The phosphorus requirements of growing cattle consuming forage diets (Ternouth *et al.*), 503

Pisum
The effect of cover crops on the establishment and yield of vining peas and on nitrate leaching (Knott), 471

Plant population density
Response of sunflower (*Helianthus annuus*) to planting pattern and population density in a desert climate (Esechie *et al.*), 455

Pollination
The role of wind and insects in cashew (*Anacardium occidentale*) pollination in NE Brazil (Freitas and Paxton), 319

Potato
The yield of potato plants as affected by stem canker (*Rhizoctonia solani*), blackleg (*Erwinia carotovora* subsp. *atroseptica*) and by neighbouring plants (Hide *et al.*), 429
General adaptability of potato genotypes selected in the UK for the Mediterranean region (Brown *et al.*), 441

Rice
The role of tolerant genotypes and plant nutrients in the management of iron toxicity in lowland rice (Sahrawat *et al.*), 143

Roots
Stimulation of root growth and soil nitrogen uptake by foliar application of urea to wheat and sunflower (Sen and Chalk), 127

Rumen
The dacron bag technique for comparing rumen degradability of untreated and ammonia-treated barley straw – effect of particle size and degree of replication (Pike *et al.*), 201

Ryegrass, *see Lolium*

Salinity
The effects of soil sodicity on emergence, growth, development and yield of oilseed rape (*Brassica napus*) (Gutiérrez Boem and Lavado), 169

Seeds
Hardseededness and pattern of hard seed breakdown in Sardinian populations of *Medicago polymorpha* under field conditions (Porqueddu *et al.*), 161

Seed yield and hardseededness of two amphicarpic pasture legumes (*Vicia sativa* ssp. *amphicarpa* and *Lathyrus ciliolatus*) and two annual medics (*Medicago rigidula* and *M. noeana*) (Christiansen *et al.*), 421

Shading

Persistence of tropical forage grasses in shaded environments (Wong and Stür), 151

Sheep

Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*) (Wang *et al.*), 87

The effect of sucrose supplementation on kinetics of nitrogen, ruminal propionate and plasma glucose in sheep (Sutoh *et al.*), 99

Effect of growth hormone on the response to insulin and glucose turnover in sheep (Rose and Obara), 107

Physical structure of twelve forage species in relation to rate of intake by sheep (Wilman *et al.*), 277

Effect of condensed tannins in *Lotus corniculatus* upon lactation performance in ewes (Wang *et al.*), 353

Pre- and post-natal wool follicle development and density in sheep of five genotypes (Hocking Edwards *et al.*), 363

Uptake of labelled phenylalanine into different blood fractions in the portal vein and cranial mesenteric vein of lambs (Neutze *et al.*), 511

Slurries

Sewage sludge applications to grassland: influence of sludge type, time and method of application on nitrate leaching and herbage yield (Misselbrook *et al.*), 343

Effects of application technique and anaerobic digestion on gaseous nitrogen loss from animal slurry applied to ryegrass (*Lolium perenne*) (Rubæk *et al.*), 481

Soils

Effect of nitrogen fertilizer applied to winter oilseed rape (*Brassica napus*) on soil mineral nitrogen after harvest and on the response of a succeeding crop of winter wheat to nitrogen fertilizer (Shepherd and Sylvester-Bradley), 63

Factors affecting nitrate leaching from a calcareous loam in East Anglia (Davies *et al.*), 75

The effects of soil sodicity on emergence, growth, development and yield of oilseed rape (*Brassica napus*) (Gutiérrez Boem and Lavado), 169

Availability to subsequent crops and leaching of nitrogen in ^{15}N-labelled sugarbeet tops and oilseed rape residues (Thomsen and Christensen), 191

Sewage sludge applications to grassland: influence of sludge type, time and method of application on nitrate leaching and herbage yield (Misselbrook *et al.*), 343

Sowing date

Grain yield, biomass and leaf area of triticale in response to sowing date and cutting stage in three contrasting Mediterranean environments (Ramos *et al.*), 253

Sowing rates

Influence of sowing rate on dry matter yield, plant density and survival of lucerne (*Medicago sativa*) under dryland and irrigated conditions (Thompson and Stout), 301

Sugarbeet

Availability to subsequent crops and leaching of nitrogen in ^{15}N-labelled sugarbeet tops and oilseed rape residues (Thomsen and Christensen), 191

Sulphur

Sulphur uptake, yield responses and the interactions between nitrogen and sulphur in winter oilseed rape (*Brassica napus*) (McGrath and Zhao), 53

Sunflower, *see Helianthus*

Tannins

Effect of condensed tannins upon the performance of lambs grazing *Lotus corniculatus* and lucerne (*Medicago sativa*) (Wang *et al.*), 87

Effect of condensed tannins in *Lotus corniculatus* upon lactation performance in ewes (Wang *et al.*), 353

Techniques

The dacron bag technique for comparing rumen degradability of untreated and ammonia-treated barley straw – effect of particle size and degree of replication (Pike *et al.*), 201

Administration in suspension-form of n-alkane external markers for dry matter intake and diet selection studies (Marais *et al.*), 207

Physiological responses of farmed red deer to management practices and their modulation by long-acting neuroleptics (Diverio *et al.*), 211

Comparison between reconstituted sheep faeces and rumen fluid inocula and between *in vitro* and *in sacco* digestibility methods as predictors of intake and *in vivo* digestibility (Nsahlai and Umunna), 235

Effects of recurrent selection on populations of various generations in wheat by using the Tai Gu single dominant male-sterile gene (Wang *et al.*), 397

Effects of application technique and anaerobic digestion on gaseous nitrogen loss from animal slurry applied to ryegrass (*Lolium perenne*) (Rubæk *et al.*), 481

Uptake of labelled phenylalanine into different blood fractions in the portal vein and cranial mesenteric vein of lambs (Neutze *et al.*), 511

Temperature

Effects of cold conditions on heat production by young sambar (*Cervus unicolor*) and red deer (*Cervus elaphus*) (Semiadi *et al.*), 221

Seed-size effects on lentil (*Lens culinaris*) yield potential and adaptation to temperature and rainfall in West Asia (Erskine), 335

Trifolium

Forage yield and quality interactions between white clover and contrasting ryegrass varieties in grazed swards (Evans *et al.*), 295

Triticale

Grain yield, biomass and leaf area of triticale in response to sowing date and cutting stage in three contrasting Mediterranean environments (Ramos *et al.*), 253

Vicia faba

Some morphological characters and yield of broad bean (*Vicia faba*) as affected by pesticide treatments (El-Sayed *et al.*), 463

Vigna

Phenological development in bambara groundnut (*Vigna subterranea*) transferred from 14 to 11 h photoperiods (Linnemann), 175

Growth, yield and plant–water relations of four cowpea

(*Vigna unguiculata*) cultivars in the Sahel (Sivakumar *et al.*), 183
Growth, development and yield of bambara groundnut (*Vigna subterranea*) in response to soil moisture (Collinson *et al.*), 307

Water relations
Growth and water relations of diverse populations of *Lolium perenne* exposed to drought in field, glasshouse and controlled environment (Amin and Thomas), 15
Growth, yield and plant–water relations of four cowpea (*Vigna unguiculata*) cultivars in the Sahel (Sivakumar *et al.*), 183
Growth, development and yield of bambara groundnut (*Vigna subterranea*) in response to soil moisture (Collinson *et al.*), 307

Wheat
Effect of nitrogen fertilizer applied to winter oilseed rape (*Brassica napus*) on soil mineral nitrogen after harvest and on the response of a succeeding crop of winter wheat to nitrogen fertilizer (Shepherd and Sylvester-Bradley), 63
Stimulation of root growth and soil nitrogen uptake by foliar application of urea to wheat and sunflower (Sen and Chalk), 127
Effects of herbicide and nitrogen fertilizer application on grain yield and quality of wheat and barley (Grundy *et al.*), 379
Rate of change in harvest index during grain-filling of wheat (Moot *et al.*), 387
Effects of recurrent selection on populations of various generations in wheat by using the Tai Gu single dominant male-sterile gene (Wang *et al.*), 397
Differences in phasic development rate amongst wheat cultivars independent of responses to photoperiod and vernalization. A viewpoint of the intrinsic earliness hypothesis (Slafer), 403

White clover, *see Trifolium*

Wool
Pre- and post-natal wool follicle development and density in sheep of five genotypes (Hocking Edwards *et al.*), 363

Yield
Grain yield, biomass and leaf area of triticale in response to sowing date and cutting stage in three contrasting Mediterranean environments (Ramos *et al.*), 253